

## Galway: GAPP - Galway Antimicrobial Prescribing Policy / Guidelines (GAPP): Bone and Joint

### Bone and Joint

1. Discuss all cases with Microbiology or Infectious Diseases.
2. Microbiological diagnosis is essential, relevant microbiological samples (bone/synovial fluid AND blood cultures) should be sent prior to treatment.
3. In patients with sepsis, antibiotic therapy should not be delayed while awaiting bone or synovial fluid sampling.
4. The regimens below may NOT cover Multi-drug Resistant Organisms (MDRO) in all cases. See note on [MDRO](#).

#### Empiric Antibiotics for Bone and Joint Infections

Infection	1 <sup>st</sup> Line Antibiotics	Penicillin allergy: delayed onset non-severe reaction  See <a href="#">penicillin hypersensitivity section</a> for further information	Penicillin allergy: immediate or severe delayed reaction	Comment
The regimens below may NOT cover Multi-drug Resistant Organisms (MRDO) in all cases. See note on <a href="#">MDRO</a>				
<b>Native joint Septic Arthritis</b> Nil risk factors for <a href="#">MRSA</a>	Flucloxacillin IV 2g every 6 hours	CeFAZolin IV 2g every 8 hours	<a href="#">Vancomycin</a> IV infusion, dose per GAPP App Calculator. See footnote* re monitoring.	Adequate drainage of joint fluid essential.  Discuss all cases with Microbiology or Infectious Diseases.
Known colonisation/risk factors for <a href="#">MRSA</a>		<a href="#">Vancomycin</a> IV infusion, dose per GAPP App calculator. See footnote* re monitoring.		
Risk of gonococcal infection  OR  High risk of Gram-negative organisms  (Elderly, Nursing home resident, recurrent UTI)	CefTRIAXone IV 2g every 24 hours	Discuss with Microbiology or Infectious Diseases		
<b>Osteomyelitis</b>	Discuss with Microbiology or Infectious Diseases; treat according to culture results.			Short course oral antibiotic therapy may be appropriate for acute or chronic osteomyelitis
<b>Prosthetic Joint Infection</b>	Discuss with Microbiology or Infectious Diseases; treat according to culture results.			

\*For advice on monitoring see [Vancomycin](#) Dosing & Monitoring section.

Ref:

BSR & BHPR, BOA, RCGP and BSAC guidelines for management of the hot swollen joint in adults [Rheumatology 2006;45:1039-1041](#)

Ravn C, Neyt J, Benito N, Abreu MA, Achermann Y, Bozhkova S, Coorevits L, Ferrari MC, Gammelsrud KW, Gerlach UJ, Giannitsioti E, Gottliebsen M, Jørgensen NP, Madjarevic T, Marais L, Menon A, Moojen DJ, Pääkkönen M, Pokorn M, Pérez-Prieto D, Renz N, Saavedra-Lozano J, Sabater-Martos M, Sendi P, Tevell S, Vogely C, Soriano A, The Sanjo Guideline Group. Guideline for management of septic arthritis in native joints (SANJO). *J Bone Jt Infect*. 2023;

Éric Senneville, Zaina Albalawi, Suzanne A van Asten, Zulfiqarali G Abbas, Geneve Allison, Javier Aragón-Sánchez, John M Embil, Lawrence A Lavery, Majdi Alhasan, Orhan Oz, İlker Uçkay, Vilma Urban■■■-Rovan, Zhang-Rong Xu, Edgar J G Peters, IWGDF/IDSA Guidelines on the Diagnosis and Treatment of Diabetes-related Foot Infections (IWGDF/IDSA 2023), Clinical Infectious Diseases, 2023;, ciad527,  
<https://doi.org/10.1093/cid/ciad527>