
Galway: GAPP - Galway Antimicrobial Prescribing Policy / Guidelines (GAPP): Continuing management of neutropenic sepsis - Algorithm
Galway: GAPP - Galway Antimicrobial Prescribing Policy / Guidelines (GAPP) - Last Updated: Nov. 7, 2024, 11:53 a.m., printed: Nov. 23, 2024, 10:41 a.m.
page 1 of 2

Neutropenic Sepsis Guidelines – CONTINUING Management



Other Infections - doses are for normal renal function

Other Infections

Treatment of Viral Infections

Local Herpes Simplex:

Oral: Valaciclovir PO 1g every 12 hours OR

IV: Aciclovir IV 5mg/kg (IBW*) every 8 hours

Herpes Zoster:

Aciclovir IV 10mg/kg (IBW*) every 8 hours, then change to Valaciclovir PO 1g every 8 hours when lesions healing Consider CMV serology & PCR

*IBW = Ideal Body Weight

Treatment of PJP

Co-Trimoxazole IV/PO 120mg/kg daily in 4 divided

OR 2nd Line: Pentamidine IV 4mg/kg once daily

PLUS steroids for severe

Treat for 14 days

Prophylaxis – if indicated See full guideline

Antiviral: ValACIclovir PO 500mg bd

PJP: Co-trimoxazole PO 960mg twice daily on Mon/Wed/Friday. See full guideline for alternatives.

tablets 300mg bd for 2 doses, then 300mg once daily OR Ambisome IV 1mg/kg (round to 50 or 100mg according to weight) once daily on Mon/Wed/Friday OR Fluconazole PO 50 to 400mg once daily

Antifungal: Posaconazole PO using

(click on image to enlarge)

Galway: GAPP - Galway Antimicrobial Prescribing Policy / Guidelines (GAPP) - Last Updated: Nov. 7, 2024, 11:53 a.m., printed: Nov. 23, 2024, 10:41 a.m.